

PRV

PATENT- OCH REGISTRERINGSVERKET
Patentavdelningen

Intyg Certificate

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.



(71) Sökande Amersham Biosciences AB, Uppsala SE
Applicant (s)

(21) Patentansökningsnummer 0300567-5
Patent application number

(86) Ingivningsdatum 2003-02-28
Date of filing

Stockholm, 2003-10-27

För Patent- och registreringsverket
For the Patent- and Registration Office

Hjördis Segerlund
Hjördis Segerlund

Avgift
Fee 170:-

**A METHOD OF GENERATING METAL
CHELATING AFFINITY LIGANDS**

Technical field

The present invention relates to a method of generating polydentate metal chelating affinity ligands. The invention also encompasses a method of generating a separation medium comprising such ligands, as well as such ligands and such a medium.

Background


In any chemical or bioprocessing industry, the need to separate and purify a product from a complex mixture is a necessary and important step in the production line. Today, there exists a wide market of methods in which industry can accomplish these goals, one of which is chromatography. Chromatography is quite well suited to a variety of uses in the field of biotechnology, since it can separate complex mixtures with great precision and also is suitable for more delicate products, such as proteins, since the conditions under which it is performed are not typically severe.

One chromatography method, which is an especially sensitive separation technique and also applicable to most types of proteins, is metal chelate affinity chromatography (MCAC), also known as immobilised metal ion adsorption chromatography (IMAC). This technique is commonly used in purification schemes together with another chromatographic step, such ion exchange chromatography (IEX) and/or hydrophobic interaction chromatography (HIC).

More specifically, IMAC utilises matrices that comprises a group capable of forming a chelate with a transition metal ion, which chelate in turn is used as the ligand in chromatography to adsorb a compound from a liquid. The binding strength in IMAC is affected predominately by the species of metal ion, the pH of the buffers and the nature of the ligand used. Since the metal ions are strongly bound to the matrix, the adsorbed protein can be eluted either by lowering the pH or by competitive elution.

In general, IMAC is useful for separation of proteins or other molecules that present an affinity for the transition metal ion of the matrix. For example, proteins will bind to the matrix upon the presence of accessible histidine, cysteine and tryptophan residues, which all exhibit an affinity for the chelated metal. With the advent of molecular biological techniques, proteins are now easily tailored or tagged with one or more histidine residues in order to increase their affinity to metal chelated ligands, and accordingly, metal chelate chromatography has more recently assumed a more important role in the purification of proteins. Huvudfoxen Kassan

Simple chelators have been suggested as ligands for IMAC, such as iminodiacetic acid (IDA). IDA, coupled to agarose supports and subsequent charged with various metals, such as Cu^{2+} , Zn^{2+} and Ni^{2+} , has been used for capture of proteins and peptides and is also available as commercial resins. More specifically, USP 4,551,271 (Hochuli, assigned to Hoffmann-La Roche Inc.) discloses a metal chelate resin which comprises IDA ligands, in the purification of interferon. The resin can be defined by the following formula:



wherein Me is Ni or Cu.

The best results are obtained with this resin if the interferon has already been partially purified. The resin can according to the specification be prepared in a known manner by treating agarose with epichlorohydrin or epibromohydrin, reacting the resulting epoxide with iminoacetic acid disodium salt and converting the product into the copper or zinc salt by washing with a copper (II) or zinc solution.

More recently, EP 87109892.7 (F. Hoffmann-La Roche AG) and its equivalent USP 4,877,830 (Döbeli et al, assigned to Hoffmann-La Roche Inc.) disclosed a tetradentate chelator known as nitrilotriacetic acid (NTA) for use with metals

that have six coordination sites. More specifically, the matrices can be described by the general formula

[carrier matrix] - spacer - NH - (CH₂)_x - CH(COOH) - N(CH₂COO⁻)₂Ni²⁺,

wherein x=2-4. The disclosed matrix is prepared by reacting an amino acid compound of the formula R - HN - (CH₂)_x - CH(NH₂) - COOH, wherein R is an amino protecting group and x is 2, 3 or 4, with bromoacetic acid in alkaline medium and subsequently, after an intermediate purification step, cleaving off the protecting group and reacting this group with an activated matrix. Accordingly, the method of preparation involves separate steps for alkylating and deprotecting the amino acid, which steps renders the method time-consuming and hence costly. In addition, the alkylation chemistry is less efficient, and after deprotection, the product is not well defined regarding rest products from neutralisation and cleavage. Following this, the material is coupled to a solid support that carries carboxyl functionalities by forming an amide bond. However, this procedure may involve disadvantages, since the media obtained presents the immobilised desired chelating ligand as well as some unreacted carboxylic groups, thus yielding a heterogeneous media. Furthermore, mono-N-protected amino acid compounds are expensive starting materials, rendering the overall method even more costly.

Finally, WO 01/81365 (Sigma-Aldrich Co.) discloses a metal chelating composition that according to the specification is capable of forming relatively stable chelates with metal ions and exhibits an improved selectivity for polyhistidine tagged proteins. According to said WO 01/81365, the linkage between the chelator and the resin is an important parameter for the selectivity, and the linkage is a neutral ether, a thioether, a selenoether or an amide. The disclosed compositions are coupled to an insoluble carrier, such as Sepharose™ according to given examples. The chromatographic media is produced in two different ways; either by a solid phase reaction directly on to the pre-activated solid support eventually used in the chromatographic media, or by a separate in solution

synthesis of the intermediate product N,N,N',N'-tetrakis (carboxymethyl)-L-cystine that is eventually coupled to the solid support.

The solid phase synthesis is carried out by adding L-cysteine to a previously epichlorhydrine activated Sepharose gel under alkaline conditions for a prolonged reaction time (18h), followed by washings. Thereafter bromoacetic acid is added, again under alkaline conditions and a prolonged reaction time (72 h), and again followed by washings, and eventually capping of remaining free amino groups present on the gel with acetic acid anhydride.

Solid phase synthesis in this way offers poor control of the reaction and potential side reactions, and thereby yields a less homogeneous product.

The alternative route, relying on in solution phase synthesis of an intermediate product starts with addition of a large excess (40 times) of glyoxylic acid to L-cystine in an alkaline borate buffer. The intermediate product was thereafter, after pH manipulations and conductivity adjustment of the reaction mixture, purified with ion exchange chromatography to give N,N,N',N'-tetrakis (carboxymethyl)-L-cystine.

Before coupling to a solid support the N,N,N',N'-tetrakis (carboxymethyl)-L-cystine has to be reduced to N,N -bis (carboxymethyl)-L-cysteine using tris (carboxyethyl) phosphine under alkaline conditions. This material can finally be used for coupling to a pre-activated solid support forming the chromatographic media. This synthetic method is elaborate and depends on a large excess of reagents to form the desired product that is eventually purified under specific chromatographic conditions, followed by reduction as an additional synthetic step, and is thereby less suited for use in large-scale production.

Accordingly, there is still a need of improved methods for synthesis of IMAC ligands as well as of methods for the immobilisation thereof to a base matrix.

Summary of the present invention

Thus, one object of the present invention is an improved method of generating polydentate metal chelating affinity ligands for subsequent coupling to a base

+46 18 6121830

5

Ink. t. Patent- och reg.verket

2003-02-28

Huvudfaxen Kassar

matrix, which method utilises cost-effective and easily available starting materials and reagents and provides high yields. This can be achieved as defined in claim 1.

Another object of the present invention is to enable a careful selection of the ligands that are immobilised onto a base matrix for use in IMAC. This can be achieved by a method of generating a separation medium comprising polydentate metal chelating affinity ligands coupled to a base matrix, wherein the coupling chemistry is well defined and easy to control.

A further object of the invention is to provide such a method, which results in a homogenous product.

Yet a further object of the invention is to provide a method as discussed above, which also allows introducing two or more functionalities, which functionalities may be of the same kind or different.

Another object of the present invention is to provide ligands for immobilised metal affinity chromatography, which ligands presents an improved handle for coupling to a base matrix and hence an improved coupling efficiency as compared to prior art ligands.

An additional object of the present invention is to provide a chromatographic medium, which in use for immobilised metal affinity chromatography (IMAC) yields a low leakage of metal ions.

Another object of the invention is to provide a method of generating a diverse library of metal chelating affinity ligands based on the same scaffold, which method can be used for ligand optimisation towards a specific application.

The objects of the invention can be achieved by one or more of the appended claims. Further objects, advantages and embodiments of the present invention will appear from the detailed description that follows.

Brief description of the drawings

Figure 1 provides a schematic illustration of the general route to produce a separation medium that comprises polydentate metal chelating affinity chromatography ligands according to the invention.

Figure 2 (a) and (b) illustrate purification of Maltose Binding Protein with (His)₆-tail (MBP-His) using IMAC separation medium prepared according to the invention. More specifically, Figure 2(a) is the chromatogram, while Figure 2(b) is an enlargement of the gradient part of said chromatogram. In Figure 2, the curve at A280 nm is denoted A, the percentage (%) of elution buffer is denoted B and the conductivity is denoted C.

Figure 3 shows SDS-PAGE analyses of fractions from IMAC purification of MBP-His. Fraction numbering as in Figure 2 chromatogram.

Figure 4 shows a test chromatogram, wherein UV 372 nm = A, Conductivity = B, Injection = C. The nickel-binding capacity of this prototype was determined to 16 µmol Ni/ml and the metal leakage to 4%.

Figure 5 shows the results of the nickel capacity test performed on a separation medium according to the invention as described in Example 5.

Definitions

The term "separation medium" is used herein for a material useful e.g. as packing of a chromatography column, and more specifically consists of one or more ligands coupled to a base matrix. Thus, the base matrix acts as a carrier, while the ligands provide functionalities that interact with target substances in chromatography.

The term "spacer" is used for a chemical entity that distances a ligand from the base matrix.

The term "ligand" means herein a chemical entity capable of binding target substances. Such target substances may be either a compound, which it is desired to isolate or remove by chromatography, or alternatively an analytical target substance.

The term "polydentate metal chelating" ligands refers to ligands with two or more donor atoms that can coordinate to, i.e. chelate, a metal simultaneously. Thus, a polydentate ligand has two or more donor atoms and occupies two or more sites in a coordination sphere.


Thus, the term "metal chelating functionalities" refers to the groups that provides donor atoms. Usually, the functionalities are distanced from each other and hence the term "ligand arm" is used for each functionality.

The term "gel" is used for a separation matrix, which is in the form of a gel.

Detailed description of the invention

Thus, in a first aspect, the present invention relates to a method of generating at least one polydentate metal chelating affinity ligand, which method comprises the steps of

(a) providing at least one scaffold defined by the general formula (I)



wherein X_1 , X_2 and X_3 irrespective of each other are sp^2 - or sp^3 -hybridised carbon atoms or heteroatoms,

X_4 is a nucleophile and

m is an integer of 0-2;

(b) providing at least one polydentate metal chelating affinity ligand arm, optionally in a form wherein the metal chelating functionalities are protected, on each scaffold by derivatisation of the nucleophilic X_4 groups of said scaffolds while retaining the cyclic structure of the scaffold;

- (c) ring-opening at the bond between the carbonyl and the sulphur of the derivatised scaffold by adding a reagent that adds one or more metal chelating affinity ligand arms to the scaffold; and, if required,
- (d) deprotecting the functionalities of the ligand arm(s) provided in step (b).

The scaffold of step (a) can be provided as a solid or, preferably, in a solvent. In the most advantageous embodiment, in formula (I), X_1 , X_2 and X_3 are carbon atoms. In alternative embodiments, one or more of X_1 , X_2 and X_3 are heteroatoms, i.e. selected from the group comprised of oxygen, sulphur, nitrogen and/or silica, provided that said heteroatom does not interfere in the subsequent use of the ligands.

In formula (I), X_4 is any suitable nucleophilic group that enables derivatisation. Thus, in an illustrative embodiment, X_4 is selected from a group such as $-OH$, $-SH$ or $-NH_2$. In an advantageous embodiment, X_4 is $-NH_2$.

As mentioned above, m can be any integer of 0-2, i.e. 0, 1, or 2. As appears from the method steps, the value of m will decide the number of atoms between the ligand arms and its attachment point to a base matrix when it is developed into a separation medium.

In an advantageous embodiment, in formula (I), m is 1 and the scaffold is homocysteine thiolactone. As the skilled person in this field will appreciate, homocysteine thiolactone can be used in pure or racemate form. Homocysteine thiolactone is commercially available, e.g. from Aldrich, catalogue no. H1, 580-2, and CAS no. 6038-19-3.

In step (b), the derivatisation is performed by adding a suitable derivatisation agent comprised of a first part, which is electrophilic and hence capable of reacting with X_4 of Formula (I), and a second part, which comprises a metal chelating functionality.

The first part of the derivatisation agent, i.e. electrophilic part, can be illustrated by $C=C$; $C-Y$, wherein Y represents for example a halogen, such as Br , I , Cl , or a mesylate, or a tosylate group; or an acid or an activated acid such as $WC=O$, wherein W is for example formed from N -hydrosuccinimide, pentafluorophenol, para-nitrophenol or isopropyl chloroformate.

In an advantageous embodiment, the derivatisation is provided by adding two derivatisation agents, each one of which comprises different or identical metal chelating functionalities, herein denoted L_1 and L_2 . In this embodiment, the electrophilic parts of the agents are preferably of the same nature in order to facilitate the derivatisation. In an alternative embodiment, more than two different or identical metal chelating functionalities are introduced by derivatisation of X_4 , preferably by use of two or more different steps, as is realised by the skilled person in this field. Accordingly, multiple functionalities are easily provided in the same, polydentate metal chelating affinity ligand.

The derivatisation agent(s) used in the present method can either comprise the metal chelating functionality in protected form, wherein the donor atoms are unavailable for reaction during the derivatisation of the scaffold, or in non-protected form. In the embodiment where the functionality is protected, said protecting group should be easy to remove in a subsequent step. Accordingly, the protecting group is either acid labile, such as an alkyl group, or base labile, such as a tert-butyl group. In one embodiment, the protecting group is a CH_2CH_3 group. Various metal chelating functionalities are known in this field, and can in principle be any electron-donating group. More specifically, the metal chelating functionalities used in the present method are selected from the group that consists of aromates, heterocyclic derivatives, such as pyridine, thiophene, furan and imidazole, acids, esters, ketones, amides, sulphones, sulphonamides, nitrile, carbon-carbon double and triple bonds.

In an illustrative embodiment, the derivatisation agent is a halogenated carboxylic acid ester, such as a halogenated carboxylic acid alkyl ester. As mentioned above, in a more specific embodiment, X_4 is NH_2 . Methods for reacting an NH_2 group with a group carrying a halogen or another leaving group are well known in this field and are conveniently performed at ambient temperature in a solvent such as N,N-dimethylformamide (DMF). In one embodiment, in order to provide two metal chelating affinity ligand arms on each scaffold by the derivatisation, the derivatisation agent is used in a molar ratio of 2:1 to the scaffold. The skilled person can easily monitor the reaction and confirm the derivatisation obtained by a conventional method such as LC-MS. Thus, the present invention provides a less complex synthetic route to polydentate metal chelating affinity ligands than WO 01/81365. Due to the advantageous chemistry, the present method also results in a more homogenous product. The yields obtained according to the present method can be as high as 90%, and the starting materials are at present readily available to a reasonable cost.

In step (c), ring-opening is provided at the bond between the carbonyl and the sulphur of the derivatised scaffold by adding a reagent, which adds one or more metal chelating affinity ligand arms of the scaffold. Thus, the cyclic structure is opened up in order to provide one or more further metal chelating ligand arms.

The ring opening will also result in a handle available for subsequent coupling to a base matrix in the form of the thiophilic group, which due to its nucleophilic nature provides convenient coupling chemistry.


In an advantageous embodiment, the ring opening is hydrolysis by addition of alkali hydroxide, such as NaOH, in which case the carbonyl of the scaffold is transformed into a carboxylic group. However, as the skilled person in this field will understand, if the ring opening is performed with a different reagent, one or more different metal chelating functionalities can be introduced. In an alternative embodiment, the ring opening is aminolysis, in which case the ni-

trogen carries one or more metal chelating functionalities. In this embodiment, the reagent is defined by the general formula II:

Ink. t. Patent- och reg.verket

2003 -02- 2 8

Huvudfaxen Kassar



Formula 2

wherein L_3 and L_4 comprise metal chelating functionalities, which can be the same or different. Further, in one embodiment, said ligand arms L_3 and L_4 are the same as those provided in step (b) by the derivatisation.

Thus, as indicated above, in the embodiment where the metal chelating functionalities L_1 and L_2 were protected during the derivatisation step, a step of deprotection is preferably performed. In one embodiment, said deprotection is performed as a separate step that follows step (c), and can be achieved by adding a base or an acid, as indicated above. The chemistry useful for protection/deprotection of functionalities are well-known in this field, and the skilled person in this field can easily perform such steps.

In an especially advantageous embodiment of the present method, the deprotection is performed in the same step as the ring-opening. Accordingly, the great advantage of this embodiment is that polydentate metal chelating affinity ligands can be generated using a two step procedure. Consequently, this embodiment provides a less complex method than the prior art methods for the synthesis of polydentate metal chelating affinity ligands. In one embodiment, wherein the derivatisation agent comprises a base labile group, this step is provided by addition of sodium hydroxide. The hydrolysis is advantageously performed at ambient temperature for e.g. 1-2 hours. In fact, the present inventors have also shown that even though all the starting material can be converted

within 90 minutes, an additional 48 hours does not give rise to any side products. Accordingly, the hydrolysis according to the invention results in a stable, homogenous and well-defined product. In an alternative embodiment, wherein the derivatisation agent comprises an acid labile group, in step (c), this step is provided by addition of an acid, such as HCl.

In a specific embodiment, in the present method, steps (a) and (b) have been performed earlier to provide a ready-derivatised scaffold. Accordingly, the present invention also encompasses a method, wherein the carboxymethylation of the scaffold has been performed earlier.

In an advantageous embodiment, the product so obtained is coupled via its sulphur to a base matrix in order to produce a separation medium. Such a separation medium is useful for isolation of target substances, for analytical purposes etc. The base matrix used in the present method can be of any material suitable for the intended use.

Thus, in the case where the separation medium is intended for use in immobilised metal chelating affinity chromatography, the base matrix is commonly in beaded or monolithic form and made from natural polymers, e.g. agarose or dextran, or synthetic polymers, such as divinylbenzene or styrene. The base matrix can e.g. be in the form of a gel.

As regards natural polymers, suitable porous polymer beads thereof are either easily performed by the skilled person in this field according to standard methods, such as inverse suspension gelation (S Hjertén: Biochim Biophys Acta 79(2), 393-398 (1964) or spinning disk technique (see e.g. WO 88/07414 (Prometic Biosciences Inc)). Alternatively, natural polymer beads are obtained from commercial sources, such as Amersham Biosciences AB, Uppsala, Sweden. Illustrative tradenames of such useful natural polymer beads are e.g. of the kind known as Sepharose™ or Sephadex™.

As regards synthetic polymers, the base matrix is comprised of cross-linked synthetic polymers, such as styrene or styrene derivatives, divinylbenzene, acrylamides, acrylate esters, methacrylate esters, vinyl esters, vinyl amides etc. Such polymers are easily produced according to standard methods, see e.g. "Styrene based polymer supports developed by suspension polymerization" (R Arshady: Chimica e L'Industria 70(9), 70-75 (1988)). Alternatively, a commercially available product, such as Source™ (Amersham Biosciences AB, Uppsala, Sweden) can be surface-modified according to the invention.

In alternative embodiments, the base matrix can e.g. be a membrane, a filter, one or more chips, surfaces, capillaries etc.

In one embodiment, the reactive groups of the base matrix are allyl groups i.e. carbon-carbon double bonds. In one embodiment, a commercially available base matrix, which already exhibits allyl groups is used. In an alternative embodiment, the allyl groups are provided according to well known methods. Thus, in an illustrative embodiment, the present base matrix has been allylated by treatment with an epoxide carrying an allyl functionality at suitable temperatures and reaction times. One example of such a commonly used allyl-functional epoxide is allyl glycidyl ether (AGE). Accordingly, in a specific embodiment, in step (d), the sulphur group of the ligand is coupled to the base matrix via the activated allyl group of allyl glycidyl ether (AGE). In this embodiment, in the final product, the sulphur group will be attached to the base matrix by a spacer comprising ether groups and hydroxy groups, and the separation medium can be defined as base matrix -O-CH₂-CHOH-CH₂-O-CH₂-CHOH-CH₂-S- ligand.

In alternative embodiments, other well-known coupling techniques of thiol-containing ligands are used, such as opening of epoxide or radical addition to double bonds.


In a specific embodiment, said allyl groups are activated by bromination or alternatively, the coupling is a free radical reaction. The free radical used can be any suitable commercially available initiator, UV etc.

A second aspect of the present invention is a polydentate metal chelating affinity ligand or a separation medium comprising at least one, preferably a plurality of, polydentate metal chelating affinity ligands coupled to a base matrix, which medium has been generated by a method as described above. In a specific embodiment, the metal chelating affinity ligands are tridentate. Such a separation medium can then be charged with a suitable metal ion, such as Cu(II), Zn(II), Ni(II), Ca(II), Co(II), Mg(II), Fe(III), Al(III), Ga(III), Sc(III) etc, and used according to well known principles of IMAC, e.g. as outlined in the section "Background" above. In the most preferred embodiment, Ni^{2+} is used.

In an advantageous embodiment, the present polydentate metal chelating affinity ligands are tridentate ligands are defined by the formula

$$-S - (\text{CH}_2)_n - \text{CH}(\text{COOH}) - \text{N}(\text{CH}_2\text{COO}^-)_2.$$

In a specific embodiment, the present separation medium, which comprises polydentate metal chelating affinity ligands coupled to a base matrix, is defined by the general formula



wherein n is an integer of 2-4. In one embodiment, $n = 2$. In this context, it is understood that if the base matrix is e.g. in the form of a particle, then a plurality of ligands will be coupled to each particle as described above.

A third aspect of the invention is the use of homocysteine thiolactone as a starting material in the preparation of polydentate metal chelating ligands. The

invention also encompasses the use of a carboxymethylated scaffold such as homocysteine thiolactone in the preparation of polydentate metal chelating ligands. In the most advantageous embodiment, said use is as defined above. As mentioned above, homocysteine thiolactone is commercially available.

A further aspect of the present invention is a kit, which comprises a scaffold as defined by the general formula (I) above, which kit comprises said scaffold in a solid state together with instructions, preferably written, for use thereof in the manufacture of metal chelating affinity ligands or a separation medium comprising polydentate metal chelating affinity ligands coupled to a base matrix. In alternative embodiment, a kit according to the invention comprises any other form of the scaffold, such as a partly or fully derivatised scaffold, together with liquids and/or reagents suitable for performing the method according to the invention. In a specific embodiment, a kit is comprised of a scaffold reacted according to the present method except the deprotection, in which case the kit also comprises a suitable reagent for deprotection, such as a base or an acid, together with instructions for use.

The present invention also encompasses a chromatography column packed with a medium according to the invention. The column can be of any size, such as for large-scale production or lab-scale, or suitable for analytical purpose. The column can also be combined with separation medium and optionally liquids into a second kind of kit, which is also encompassed by the present invention. In one embodiment, the kit according to the invention comprises metal ions, such as Ni^{2+} ions.

In addition, the present invention also relates to a process of separating a target substance from a liquid, which process comprises to provide a separation medium as defined above, to charge said medium with suitable metal ions to form chelates and to contact said medium with the liquid to adsorb the target substance thereon. In an advantageous embodiment, the process also comprises a

step of eluting the target substance from the separation medium by adding a liquid that desorbs the target compound from the separation medium. In one embodiment, the elution is obtained by using a liquid that comprises a decreasing pH gradient or by applying a gradient giving an increasing imidazole concentration. The general principles of chromatography for separating a target substance as discussed above are well-known in this field, and the skilled person in this field can easily adopt the necessary parameters for use of the present process.

Finally, the present invention relates to a procedure of generating one or more diverse libraries of metal chelating affinity ligands for screening and optimising purposes. Thus, in this procedure, one arm carrying a dentate can be kept constant while other arms are selected in terms of optimal performance. As the skilled person will understand, for example one or more of the above-discussed L_1 , L_2 , L_3 and L_4 can be varied in order to identify the optimal form, and subsequently, once the optimised form has been identified, it is kept constant while other(s) are varied. Accordingly, the procedure of optimisation provides a tool to manufacture a separation medium comprising optimal, selected ligands.

Detailed description of the drawings

Figure 1 provides a schematic illustration of the general route to produce a separation medium that comprises polydentate metal chelating affinity chromatography ligands according to the invention. The first step corresponds to step (b) of the present method, i.e. a derivatisation, the second step is a hydrolysis to open up the ring of the derivatised scaffold and the last step is immobilisation i.e. coupling of the ligand so produced to a base matrix. As appears from the above, the second step is advantageously a combined ring-opening and deprotection. In Figure 1, R denotes either hydrogen or alternatively an acid or base labile protecting group.

Figure 2 (a) and (b) illustrate purification of Maltose Binding Protein with (His)₆-tail (MBP-His) using IMAC separation medium according to the inven-

tion. More specifically, Figure 2(a) is the chromatogram, while Figure 2(b) is an enlargement of the gradient part of said chromatogram. In Figure 2, the curve at A280 nm is denoted A, the percentage (%) of elution buffer is denoted B, and the conductivity is denoted C.

Figure 3 shows SDS-PAGE analyses of fractions from IMAC purification of MBP-His. Fraction numbering as in Figure 2 chromatogram.

Figure 4 shows a test chromatogram, wherein UV 372 nm = A, Conductivity = B, Injection = C. The nickel-binding capacity of this prototype was determined to 16 μ mol Ni/ml and the metal leakage to 4%, as explained in example 5 below.

Figure 5 shows the results of the nickel- capacity test performed on a separation medium according to the invention as described in Example 5. More specifically, the X axis shows the ligand density while the Y axis shows the nickel-binding. It appears clearly that the nickel-binding capacity increases linearly with the ligand density and that the slope is close to one.

EXPERIMENTAL PART

The following examples are provided for illustrative purposes only and should not be interpreted to limit the scope of the present invention as defined by the appended claims. All references given below or elsewhere in the present application are hereby included herein by reference.

Experimental

$^1\text{H-NMR}$, $^{13}\text{C-NMR}$, CH-correlation, APT and cosy spectra were recorded in δ scale (ppm) with Bruker 300 MHz using TMS as reference. All spectra were recorded in CDCl_3 unless otherwise stated. TLC was carried out using Merck precoated silica gel F₂₅₄ plates. Ninhydrin or a mixture of Mo/Ce was used to visualise spots on TLC plates. LC-MS data were recorded using Hewlett Packard 1100 MSD electro spray. The flash column chromatographic purifications were carried out using Merck G-60 silica gel.

Example 1: Carboxymethylation of scaffold to provide N,N-bis(ethyl-carboxymethyl ester)+/-homocysteine thiolactone

In a dry 250mL round-bottomed flask, the D/L homocysteine thiolactone (4.5g, 29.22 mmol) was dissolved in 100mL DMF. To this was added bromo-acetic acid ethyl ester (9.76 g, 58.44 mmol, 6.48 mL), KI (4.850 g, 29.22mmol) and NaHCO₃ (14.727g, 175 mmol). The reaction mixture was stirred at ambient temperature. The reaction was followed by TLC (toluene: ethyl acetate 3: 1) and LC-MS data. The reaction was completed after 3.5h.

The desired product has a R_f = 0.35 (toluene: ethyl acetate 3: 1). The solvent was evaporated, the resulting solid redissolved in CHCl₃, extracted with H₂O X 2. The organic phase was finally dried over Na₂SO₄, filtered and evaporated. The product was purified on flash column chromatography (toluene:ethyl acetate 3:1). Yield:7.636g (26.422mmol), 90%.

¹H-NMR : δ 1.52 (t, 6H, CH₃-CH₂-O-), δ 2.05-2.56 (m, 2H, -S-CH₂-CH₂-CH-C=O), δ 3.25 (m, 2H, -S-CH₂-CH₂-CH-C=O), δ 3.52 (s, 4H, -N-CH₂-N-CH₂-), δ 3.65 (dd, 1H, S-CH₂-CH₂-CH-C=O), δ 4.24 (m, 4H, CH₃-CH₂-O-); ¹³C-NMR : δ 14.71 (CH₃-CH₂-O-), δ 27.42 (S-CH₂-CH₂-CH-C=O), δ 29.68 (S-CH₂-CH₂-CH-C=O), δ 54.01 (CH₃-CH₂-O-), δ 60.99 (-N-CH₂-N-CH₂-), δ 67.15 (-S-CH₂-CH₂-CH-C=O), δ 170.94 (O=C-CH₂-N-CH₂-C=O), δ 207.20 (-S-CH₂-CH₂-CH-C=O). LC-MS : M⁺ 290.

Example 2: Hydrolysis to provide N,N-bis(carboxymethyl)+/-homocysteine and stability test

N,N-bis(ethyl carboxymethyl ester)+/-homocysteine thiolactone (50 mg, 0.173 mmol) prepared as described in example 1 above, was dissolved in 1mL 1M NaOH.

In a 100mL round-bottomed flask. The reaction mixture was stirred at ambient temperature for 100 minutes. The reaction was followed to completion until no starting material was visible according to LC-MS.

Stability test : After complete hydrolysis, the above reaction mixture was diluted to 5mL with H₂O. The pH was adjusted to 12.5 and the reaction mixture was heated at 50°C with stirring. A 50µL sample was withdrawn from the mixture with 1h intervals for four hours. Each 50µL sample was then mixed with 1mL MeOH for the LC-MS analysis. After this step, the reaction mixture was allowed to stand overnight and finally a 50µL sample was withdrawn for the LC-MS analysis as described above. No decomposition was observed during the time of the experiment.

The crude product was freeze-dried.

¹H-NMR (D₂O): δ 1.52 (m, 2H, -CH-CH₂-CH₂-SH), δ 2.24 (m, 2H, -CH-CH₂-CH₂-SH), δ 3.20 (m, 5H, O=C-CH₂-N-CH₂-); ¹³C-NMR (D₂O) : δ 22.34 (-CH-CH₂-CH₂-SH), δ 35.39 (-CH-CH₂-CH₂-SH), δ 57.33 (O=C-CH₂-N-CH₂-), δ 67.30 (-CH-CH₂-CH₂-SH), δ 180.55 (O=C-CH₂-N-CH₂-C=O), δ 181.84 (-N-CH-C=O); LC-MS : M⁺ 252.

Example 3: Coupling of the product obtained from example 2 to allylated agarose using AGE

10 ml Sepharose™ HP-allyl (Amersham Biosciences, Uppsala, Sweden) (44 µmol/ml gel) was stirred in 20 ml distilled water with 1g NaOAc. Br₂ aq. saturated was added until that a persisting yellow colour is formed. Sodium formate was then added until the yellow colour has disappeared. The gel was then washed with distilled water. N,N-bis(ethyl carboxymethyl ester)+/- homocysteine thiolactone (102 mg) was stirred in 2 ml 1M NaOH at room temperature for 2 hours. 5 ml distilled water and 3 ml 1M NaHCO₃ was added, and the pH was adjusted to 11.0 with 2M NaOH.

The ligand solution was then added to the drained gel in a vial, which was capped.

The vial was shaken at 50°C for 16 h, then the gel was washed with distilled water on a glass filter funnel.

Example 4: Purification of Maltose Binding Protein with (His)₆-tail (MBP-His) using an IMAC separation medium generated according to the invention and metal leakage test

Materials and methods:

Extract with MBP-His

C-terminally hexaHis-tailed Maltose Binding Protein, MBP-His, theoretical Mr and pI was 43 781 and 5.4:

Fermentation of an *E. coli* clone expressing MBP-His and cell homogenisation was performed according to a standard procedure. The concentration of MBP-His in this extract was estimated to approximately 1.9 mg/ml.

IMAC A-buffer:

For one litre: (one PBS tablet to 1000 ml water is specified to give 10 mM Na-phosphate, 140 mM NaCl and 3 mM KCl, pH 7.4.) Two PBS tablets were dissolved in water, NaCl was added from 5 M stock to give an extra 720 mM (thus finally 140+140+720 mM = 1 M), pH adjusted to 7.4 with NaOH and final volume to 1000 ml. (Buffer also contains 6 mM KCl).

Elution Buffer (IMAC B-buffer):

Prepared in the same way as IMAC A-buffer, but also imidazole was added to 500 mM (from 2.0 M stock of imidazole-HCl, pH 7.4) before final adjustment of pH and volume.

Ni²⁺-sulphate solution: 100 mM in water. Filter 0.2 µm. pH ≈ 4.6.

Sample buffer and running conditions according to Instructions for ExcelGel™ (Amersham Biosciences, Uppsala, Sweden).

Gels: ExcelGel™ SDS, gradient 8-18 %.

Sample prep: Samples were mixed 1+1 with 2 x sample buffer (for purifications with large amounts of target protein) or 1 volume sample + 1/3 volume 4x sample buffer (purifications with small amounts of target protein). (4x sample buffer = 100 mM Tris-HAc pH 7.5, 2 % SDS.) Heating at 95°C for 3-5 minutes.

Sample application: IEF sample application pieces (5x10 mm filter paper pieces, # 80-1129-46 (Amersham Biosciences, Uppsala, Sweden) were used. Up to 30-32 paper pieces were positioned with their short sides towards the running direction. 20 µl sample cocktail was applied to each piece. The pieces were left on the gel during the whole electrophoresis.

Electrophoresis: In a Multiphor™ II (Amersham Biosciences, Uppsala, Sweden) apparatus at 15° C (circulating cooling bath). EPS 3500 Power Supply. Limiting settings: 600 V, 50 mA, 30 W.

Staining: In 0.1 % Coomassie R 350 dissolved in 30 % MeOH, 10 % HAc. Destaining in 25 % EtOH, 8% HAc.

Chromatography: The media was packed in HR 5/5 columns (Amersham Biosciences, Uppsala, Sweden) to a bed height of 5 cm = 1.0 ml bed. The media was charged with Ni²⁺ prior to use by pumping the Ni-sulphate solution onto the column (5 column volumes), followed by water and by equilibration with binding buffer (= IMAC A-buffer) with 5 mM imidazole added). A short blank run was performed by applying elution buffer, again followed by equilibration with binding buffer.

The *E. coli* extract with MBP-His, 1.0 M NaCl, 5 mM imidazole and 1 mM Huvudfaxen Kasean PMSF (freshly added) was clarified by centrifugation and by 0.45 μ m filtration. The required volume was then introduced into a Superloop™ (Amersham Biosciences, Uppsala, Sweden) through a 0.2 μ m filter shortly prior to IMAC start. 3 ml was then applied to the column. After extensive washing with binding buffer, a 20-ml linear gradient was run (to 40 % Elution buffer = 200 mM imidazole) using a ÄKTA™ Explorer 10 system (Amersham Biosciences, Uppsala, Sweden). Finally, a 5-ml push with 100 % Elution buffer (500 mM imidazole) was applied. All steps were at 1.0 ml/min.

Example 5: Determination of nickel-binding capacity and metal-leakage at pH 4.0

Nickel binding capacity

The nickel (Ni^{2+}) binding capacity of a separation medium according to the invention was determined chromatographically using an ÄKTA™ Explorer 10 system (Amersham Biosciences, Uppsala, Sweden) equipped with a UV/Vis DAD detector. The test was performed on gel to which N,N-bis(carboxymethyl)+/-homocysteine had been coupled according to earlier description, and packed in 1-ml HR5/5 columns (Figure 4). This result was compared with the ligand density present on the gel as determined by elemental analysis (nitrogen analysis, each ligand containing one nitrogen atom). The correlation between nickel-binding capacity and ligand density was considered to be very good, i.e. one ligand is carrying one metal ion. Typical results are shown in Figure 5.

Method description

A NiSO_4 solution was injected to load the gel with Ni^{2+} ions. Excess metal was removed by washing with water and phosphate buffer (20 mM PO_4 , 500 mM NaCl, pH 7.4). The nickel ions bound to the gel were eluted with EDTA, which is a very strong chelator and efficiently strips the metal ions from the gel. The peak area of the green coloured Ni-EDTA complex eluted was measured at 372

nm. A linear calibration curve was established from solutions with different concentrations of Ni-EDTA and was used for quantification. The nickel-binding capacity (Figure 4, peak 1) was given as $\mu\text{mol Ni/ml}$ packed gel.


Metal leakage

A metal leakage test was performed to test the stability of the ligand-nickel complex. The nickel-loaded gel was then washed with acetate buffer, pH 4.0, before determining the Ni-content on the gel (Figure 4).

The metal leakage was considered to be very low for all tested gels prepared according to the present invention; typically the leakage was 4 %.


Method description

Leakage of nickel ions was tested analogously to the nickel-binding capacity. However, before eluting nickel ions with EDTA, the gel was washed with ten column volumes of acetate buffer, 100 mM, pH 4.0. The leakage was given as % of the binding capacity and was determined as the difference between the nickel-binding capacity and the amount eluted after washing with pH 4 (Figure. 4, peak 2) according to the formula: $(\text{area}_{\text{Peak 2}} - \text{area}_{\text{Peak 1}}) / \text{area}_{\text{Peak 1}}$.



CLAIMS

1. A method of generating at least one polydentate metal chelating affinity ligand, which method comprises the steps of
 - (a) providing at least one scaffold defined by the general formula (I)



wherein X_1 , X_2 and X_3 irrespective of each other are sp^2 - or sp^3 -hybridised carbon atoms or heteroatoms,

X_4 is a nucleophile and

m is an integer of 0-2;

- (b) providing at least one polydentate metal chelating affinity ligand arm, optionally in a form wherein the metal chelating functionalities are protected, on each scaffold by derivatisation of the nucleophilic X_4 groups of said scaffolds, while retaining the cyclic structure of the scaffold;
 - (c) ring-opening at the bond between the carbonyl and the sulphur of the derivatised scaffold by adding a reagent that adds one or more metal chelating affinity ligand arms to the scaffold; and, if required,
 - (d) deprotecting the functionalities of the ligand arm(s) provided in step (b).
 2. A method according to claim 1, wherein in formula (I), X_1 , X_2 and X_3 are carbon atoms.
 3. A method according to any one of the preceding claims, wherein in formula (I), X_4 is $-NH_2$.
 4. A method according to any one of the preceding claims, wherein in formula (I), m is 1 and the scaffold is homocysteine thiolactone.

5. A method according to any one of the preceding claims, wherein in step (b), the derivatisation is provided by adding at least one derivatisation agent comprised of one part, which is electrophilic and hence capable of reacting with X_4 in Formula (I), and one part, which is a metal chelating affinity ligand.
6. A method according to claim 5, wherein the derivatisation is provided by adding two derivatisation agents, which comprise two different or identical metal chelating functionalities.
7. A method according to claim 5 or 6, wherein one derivatisation agent is a halogenated, protected ester.
8. A method according to claim 7, wherein one derivatisation agent is bromoacetic acid ethyl ester.
9. A method according to any one of the preceding claims, wherein in step (c), the ring-opening is hydrolysis provided by adding a base.
10. A method according to any one of the preceding claims, wherein the metal chelating functionalities are protected in step (b), and subsequently, step (c) and step (d) are performed simultaneously.
11. A method according to any one of the preceding claims, wherein steps (a) and (b) have been performed earlier to provide a ready-derivatised scaffold.
12. A method according to any one of the preceding claims, wherein the product obtained from step (d) is coupled via its thiol group to a base matrix in order to produce a separation medium.
13. A method according to claim 12, wherein the thiol group is coupled to allyl groups of the base matrix.
14. A method according to claim 12 or 13, which also includes a step of allylation of a base matrix to provide reactive groups.
15. A method according to any one of the claims 12-14, wherein the thiol group of the ligand is coupled to the base matrix via the allyl group of allyl glycidyl ether (AGE).
16. A method according to any one of the claims 12-15, which also comprises a step of activating the reactive groups of the base matrix.

17. A method according to claim 16, wherein the activation is performed by bromination.
18. A polydentate metal chelating affinity ligand, or a separation medium comprising a plurality of polydentate metal chelating affinity ligands coupled to a base matrix, which ligand or medium has been generated by a method according to any one of claims 1-17.
19. A ligand or separation medium according to claim 18, wherein the ligand is a tridentate ligand.
20. Use of homocysteine thiolactone as a starting material in the preparation of polydentate metal chelating affinity ligands.
21. A kit which comprises a scaffold as defined by the general formula (I) as defined in any one of claims 1-4, which kit comprises said scaffold in a solid state together with written instructions for use thereof in the manufacture of a separation medium comprising a plurality of polydentate metal chelating affinity ligands coupled to a base matrix.
22. A kit according to claim 21, wherein the scaffold is homocysteine thiolactone.
23. A separation medium comprising polydentate metal chelating affinity ligands coupled to a base matrix and defined by the general formula
base matrix-O-CH₂-CHOH-CH₂-O-CH₂-CHOH-CH₂S - (CH₂)_n - CH(COOH)
- N(CH₂COO⁻)₂Ni²⁺
wherein n is an integer of 2-4.
24. A medium according to claim 23, wherein n = 2.
25. A chromatography column for immobilised metal ion adsorption chromatography (IMAC), which column has been packed with a separation medium as defined in any one of claims 18-19 or 23-24.
26. A process of separating a target substance from a liquid, which process comprises to provide a separation medium according to any one of claims 18-19 or 23-24, to charge said medium with suitable metal ions to form chelates and to contact said medium with the liquid to adsorb the target substance thereon.

27. A process according to claim 26, which also comprises a step of eluting the target substance from the separation medium by adding a liquid that desorbs the target compound from the separation medium.
28. A process according to claim 27, wherein the elution is obtained by use of a liquid that comprises a decreasing pH gradient or by applying a gradient of increasing imidazole concentration.

+46 18 6121830

28

Ink. t. Patent- och reg.verket

2003 -02- 2 8

Huvudfaxen Kassan

ABSTRACT

The present invention relates to a method of generating at least one polydentate metal chelating affinity ligand, which method comprises the steps of

- (a) providing at least one cyclic scaffold comprising a carbonyl, an adjacent sulphur and a nucleophile;
- (b) providing at least one polydentate metal chelating affinity ligand arm, optionally in a form wherein the metal chelating functionalities are protected, on each scaffold by derivatisation of the nucleophile of said scaffolds, while retaining the cyclic structure of the scaffold;
- (c) ring-opening at the bond between the carbonyl and the sulphur of the derivatised scaffold by adding a reagent which adds one or more metal chelating affinity ligand arms to the scaffold; and, if required,
- (d) deprotecting the functionalities of the at least one ligand arm provided in step (b).

In the most preferred embodiment of the method, steps (c) and (d) are performed simultaneously as one single step.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131




Fig. 1

Ink. t. Patent- och reg.verket

2003 -02- 2 8

Huvudfaxen Kassar




Fig. 2

Ink. t. Patent- och reg.verket

2003 -U2- 2 8

Huvudfaxen Kassan




Fig. 3




Fig. 4

Ink. t. Patent- och reg.verket

2003-02-28

Huvudfaxen Kasse




Fig. 5